



## TREATMENT TRACKER

# Tips to Start and Stay on Your TUKYSA Treatment Plan



TUKYSA is a prescription medicine used with the medicines trastuzumab and capecitabine to treat adults with human epidermal growth factor receptor-2 (HER2) positive breast cancer that has spread to other parts of the body such as the brain (metastatic), or that cannot be removed by surgery, and who have received one or more anti-HER2 breast cancer treatments.

It is not known if TUKYSA is safe and effective in children.

**Please see Important Safety Information throughout and accompanying Important Facts about TUKYSA.**

 **TUKYSA™**  
tucatinib  
50 mg | 150 mg tablets

**SELECT SAFETY INFORMATION**

**What are the possible side effects of TUKYSA?**

TUKYSA may cause serious side effects, including:

- **Diarrhea** (watery, loose, or frequent stools) is common and can sometimes be severe. Tell your healthcare provider if you have a change in your bowel movements or severe diarrhea. Severe diarrhea can cause a loss of too much body fluid (dehydration), low blood pressure, kidney problems, and death. Your healthcare provider may prescribe medicines to treat your diarrhea during treatment with TUKYSA.


Please see Important Safety Information throughout and accompanying Important Facts about TUKYSA.



**Getting Started**

TUKYSA is part of a treatment plan that includes HERCEPTIN® (trastuzumab) and XELODA® (capecitabine).

TUKYSA is taken orally, twice-a-day



- The recommended dose of TUKYSA is 300 mg twice daily, taken 12 hours apart with or without food
- Swallow tablets whole. Do not chew, crush, or split the tablets. Do not take TUKYSA tablets if they are broken, cracked, or damaged
- If you vomit or miss a dose of TUKYSA, take your next dose at your regular time
- Your doctor may change your dose of TUKYSA if needed

- Receive trastuzumab at your healthcare provider's office or infusion center on Day 1 and again every 21 days

- Take within 30 minutes after a meal
- TUKYSA and capecitabine can be taken at the same time
- Take capecitabine for 14 days, with a 7-day break before starting again

**Remember, even though you'll stop capecitabine for the last 7 days of the schedule, TUKYSA should be taken every day.**

# My Treatment Plan—Example

Fill out the blank treatment plan found at the end of this guide with your healthcare provider or pharmacist.

**TIP!** Set AM/PM alarms to remind you to take your medications on time

## My TUKYSA™ (tucatinib) tablets Treatment Plan



TUKYSA start date 3/5

My dose is 300 mg  50-mg tablets  2 150-mg tablets 

Taken 2 times each day at 7:00 AM 7:00 PM



### My trastuzumab schedule

My infusion is given on (date) 3/5

My next infusion will be given on (date) 3/26



### My capecitabine schedule

I start on (date) 3/5

I stop on (date) 3/19

I restart on (date) 3/26

My dose is 1000 mg

Taken 2 times each day at 7:00 AM 7:00 PM

**Stay on track. Fill in your own calendar with your treatment schedule. Blank calendars can be found on pages 12 and 13. They can also be downloaded at [TUKYSA.com](http://TUKYSA.com).**

# My Treatment Calendar—Example

**Customize with your dates to keep you on track**

• Fill in the date for each day and number of each week

• Take TUKYSA every day

• Circle the days you receive trastuzumab

• Mark the days that you take capecitabine and cross out the days that you don't

• Tell your doctor if you have any side effects that bother you or do not go away

	TUKYSA Capecitabine	TUKYSA Capecitabine	TUKYSA Capecitabine	TUKYSA Capecitabine	TUKYSA Capecitabine	TUKYSA Capecitabine	TUKYSA Capecitabine
	SUN	MON	TUES	WED	THUR	FRI	SAT
DATE					3/5	3/6	3/7
AM	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
PM	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
DATE	3/8	3/9	3/10	3/11	3/12	3/13	3/14
AM	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
PM	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
DATE	3/15	3/16	3/17	3/18	3/19	3/20	3/21
AM	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
PM	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
DATE	3/22	3/23	3/24	3/25	3/26	3/27	3/28
AM	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
PM	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
DATE	3/29	3/30	3/31	4/1	4/2	4/3	4/4
AM	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
PM	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
DATE	4/5	4/6	4/7	4/8	4/9	4/10	4/11
AM	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
PM	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>



**Take notes about how you feel**

WEEK 1

*missed a dose Friday night*

WEEK 2

*Felt more tired than usual this week*

WEEK 3

WEEK 4

*minor hand pain for 2 days*

WEEK 5

WEEK 6

## Important Safety Information



### What are the possible side effects of TUKYSA?

TUKYSA may cause serious side effects, including:

- **Diarrhea** (watery, loose, or frequent stools) is common and can sometimes be severe. Tell your healthcare provider if you have a change in your bowel movements or severe diarrhea. Severe diarrhea can cause a loss of too much body fluid (dehydration), low blood pressure, kidney problems, and death. Your healthcare provider may prescribe medicines to treat your diarrhea during treatment with TUKYSA.
- **Liver Problems**, including severe cases. Your healthcare provider will test your blood to check your liver function before starting and every 3 weeks during treatment with TUKYSA, or as needed. Tell your healthcare provider right away if you have any signs and symptoms of liver problems including itching, yellowing of the skin or eyes, dark or brown urine (tea-colored), pain or discomfort in the right upper stomach area (abdomen), feeling very tired, decreased appetite, or bleeding or bruising more easily than normal.

### The most common side effects of TUKYSA:

- diarrhea
- rash, redness, pain, swelling, or blisters on the palms of your hands or soles of your feet
- nausea
- tiredness
- increased liver function blood tests
- vomiting
- mouth sores (stomatitis)
- decreased appetite
- stomach-area (abdomen) pain
- headache
- a low number of red blood cells (anemia)
- rash

**Your healthcare provider may change your dose of TUKYSA, temporarily stop, or permanently stop treatment with TUKYSA if you have certain side effects.**

TUKYSA may cause fertility problems in males and females, which may affect the ability to have children. Talk to your healthcare provider if you have concerns about fertility.

These are not all the possible side effects of TUKYSA. Discuss side effects with your healthcare provider. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/Safety/MedWatch](http://www.fda.gov/Safety/MedWatch) or call 1-800-FDA-1088.



## Important Safety Information (cont.)



### What should I tell my healthcare provider before taking TUKYSA?

Before taking TUKYSA, tell your healthcare provider about all of your medical conditions, including if you:

- have liver problems.
- are pregnant or plan to get pregnant. TUKYSA can harm your unborn baby.

#### Females who are able to become pregnant:

Your healthcare provider will do a pregnancy test before you start taking TUKYSA. Use effective birth control (contraception) during TUKYSA treatment and for at least 1 week after the last dose of TUKYSA. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with TUKYSA.

#### Males with a female partner who can get pregnant:

Use effective birth control during TUKYSA treatment and for at least 1 week after the last dose of TUKYSA.

- are breastfeeding (nursing) or plan to breastfeed. Do not breastfeed during treatment with TUKYSA and for at least 1 week after the last dose of TUKYSA.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. TUKYSA may affect the way your other medicines work, and other medicines may affect the way TUKYSA works. Keep a list of all the medicines you take and show it to your healthcare provider and pharmacist every time you get a new medicine.

### INDICATION



#### What is TUKYSA?

TUKYSA is a prescription medicine used with the medicines trastuzumab and capecitabine to treat adults with human epidermal growth factor receptor-2 (HER2) positive breast cancer that has spread to other parts of the body such as the brain (metastatic), or that cannot be removed by surgery, and who have received one or more anti-HER2 breast cancer treatments.

It is not known if TUKYSA is safe and effective in children.



## Making Your Plan

1. Fill out your treatment plan
2. Keep track of your daily dosing via the calendar
3. Remember to update the treatment plan if your healthcare provider makes changes

**Call your healthcare provider if you have any questions about TUKYSA or your treatment plan.**



For more resources  
and additional calendars,  
visit [TUKYSA.com](https://www.tukyasa.com).

Please see Important Safety Information throughout and  
accompanying Important Facts about TUKYSA.

 **SeattleGenetics**<sup>®</sup>

TUKYSA and its logo are trademarks of Seattle Genetics, Inc. Seattle Genetics and  are US registered trademarks of Seattle Genetics, Inc.  
All other trademarks are the marks of their respective owners.

© 2020 Seattle Genetics, Inc., Bothell, WA 98021 All rights reserved US-TUP-20-112-MT

 **TUKYSA**<sup>™</sup>  
tucatinib  
50 mg | 150 mg tablets

## Starting and Staying on Treatment

Here, you'll find blank calendars that you can fill in to track your treatment plan. To use:

- Fill in the date for each day
- Take TUKYSA every day
- Circle the days you receive trastuzumab
- Mark the days you take capecitabine and cross out the days you don't

Blank calendars and more resources can be found on [TUKYSA.com](https://www.tukyasa.com).

# My Treatment Plan

Fill out this blank treatment plan with your healthcare provider or pharmacist.

**TIP!** Set AM/PM alarms to remind you to take your medications on time

## My TUKYSA™ (tucatinib) tablets Treatment Plan



TUKYSA start date \_\_\_\_\_

My dose is \_\_\_\_\_ mg

\_\_\_\_\_ 50-mg tablets



\_\_\_\_\_ 150-mg tablets



Taken 2 times each day at \_\_\_\_\_ AM \_\_\_\_\_ PM



### My trastuzumab schedule

My infusion is given on (date) \_\_\_\_\_

My next infusion will be given on (date) \_\_\_\_\_



### My capecitabine schedule

I start on (date) \_\_\_\_\_

I stop on (date) \_\_\_\_\_

I restart on (date) \_\_\_\_\_

My dose is \_\_\_\_\_ mg

Taken 2 times each day at \_\_\_\_\_ AM \_\_\_\_\_ PM

Stay on track. Fill in your own calendar with your treatment schedule. Blank calendars can be downloaded at [TUKYSA.com](https://www.tukyasa.com).



# My Treatment Calendar



## Customize with your dates to keep you on track

- Fill in the date for each day and number of each week
- Take TUKYSA every day
- Circle the days you receive trastuzumab
- Mark the days that you take capecitabine and cross out the days that you don't
- Tell your doctor if you have any side effects that bother you or do not go away

	TUKYSA Capecitabine	TUKYSA Capecitabine	TUKYSA Capecitabine	TUKYSA Capecitabine	TUKYSA Capecitabine	TUKYSA Capecitabine	TUKYSA Capecitabine
	SUN	MON	TUES	WED	THUR	FRI	SAT
DATE							
AM	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
PM	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
DATE							
AM	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
PM	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
DATE							
AM	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
PM	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
DATE							
AM	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
PM	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
DATE							
AM	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
PM	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

## Take notes about how you feel

WEEK \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

WEEK \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

WEEK \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

WEEK \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

WEEK \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_



## IMPORTANT FACTS

This is only a brief summary of important information about TUKYSA. Talk to your healthcare provider or pharmacist to learn more.

## ABOUT TUKYSA

TUKYSA is a prescription medicine used with the medicines trastuzumab and capecitabine to treat adults with human epidermal growth factor receptor-2 (HER2) positive breast cancer that has spread to other parts of the body such as the brain (metastatic), or that cannot be removed by surgery, and who have received one or more anti-HER2 breast cancer treatments.

It is not known if TUKYSA is safe and effective in children.

**Important information:** Also read the capecitabine Medication Guide.

## BEFORE TAKING TUKYSA

**Tell your healthcare provider about all your medical conditions, including if you:**

- Have liver problems
- Are pregnant or plan to become pregnant. TUKYSA can harm your unborn baby

### Women who can become pregnant:

- Your healthcare provider will do a pregnancy test before you start treatment with TUKYSA
- Use effective birth control (contraception) during treatment with TUKYSA and for at least 1 week after the last dose of TUKYSA. Talk to your healthcare provider about birth control methods that you can use during this time
- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with TUKYSA

### Men with women partners who can become pregnant:

- Use effective birth control (contraception) during treatment with TUKYSA and for at least 1 week after the last dose of TUKYSA
- Are breastfeeding or plan to breastfeed. Do not breastfeed during treatment with TUKYSA and for at least 1 week after the last dose of TUKYSA

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements.

- TUKYSA may affect the way your other medicines work, and other medicines may affect the way TUKYSA works
- Keep a list of all the medicines you take and show it to your healthcare provider and pharmacist every time you get a new medicine

## HOW TO TAKE TUKYSA

- Take TUKYSA 2 times a day with or without food
- Take TUKYSA about 12 hours apart or at the same times every day
- Swallow TUKYSA tablets whole. Do not chew, crush, or split the tablets. Do not take TUKYSA tablets if they are broken, cracked, or damaged
- If you vomit or miss a dose of TUKYSA, take your next dose at your regular time

## POSSIBLE SIDE EFFECTS OF TUKYSA

**TUKYSA can cause serious side effects, including:**

- **Diarrhea** (watery, loose, or frequent stools), which is common and can sometimes be severe
  - Severe diarrhea can lead to loss of too much body fluid (dehydration), low blood pressure, kidney problems, and death
  - Tell your healthcare provider if you have a change in your bowel movements or severe diarrhea. Your healthcare provider may prescribe medicines to treat your diarrhea during treatment with TUKYSA
- **Liver problems**, including severe cases. Your healthcare provider will test your blood to check your liver function before starting TUKYSA and every 3 weeks during treatment with TUKYSA, or as needed.

Tell your healthcare provider right away if you have signs or symptoms of liver problems including itching, yellowing of the skin or eyes, dark or brown (tea-colored) urine, pain in the upper right side of your stomach area (abdomen), feeling very tired, decreased appetite, or bleeding or bruising more easily than normal.

**The most common side effects of TUKYSA include:**

- diarrhea
- rash, redness, pain, swelling, or blisters on the palms of your hands or soles of your feet
- nausea
- tiredness
- increased liver function blood tests
- vomiting
- mouth sores (stomatitis)
- decreased appetite
- stomach-area (abdomen) pain
- headache
- low red blood cell counts (anemia)
- rash

**Your healthcare provider may change your dose of TUKYSA, temporarily stop, or permanently stop treatment with TUKYSA if you have certain side effects.**

TUKYSA may cause fertility problems in males and females, which may affect the ability to have children. Talk to your healthcare provider if you have concerns about fertility.

These are not all the possible side effects of TUKYSA. Call your doctor for medical advice about side effects.

## GET MORE INFORMATION

- This is only a brief summary of important information about TUKYSA. Talk to your healthcare provider or pharmacist to learn more
- Go to [www.TUKYSA.com](http://www.TUKYSA.com) or call 1-855-4SEAGEN
- If you need help paying for your medicine, visit [www.SeaGenSecure.com](http://www.SeaGenSecure.com) for program information

